

Open Research Online

The Open University's repository of research publications and other research outputs

The spectra of lifted digraphs

Journal Item

How to cite:

Dalfó, C.; Fiol, M. A. and Širáň, J. (2019). The spectra of lifted digraphs. *Journal of Algebraic Combinatorics*, 50(4) pp. 419–426.

For guidance on citations see [FAQs](#).

© 2019 Springer Science+Business Media



<https://creativecommons.org/licenses/by-nc-nd/4.0/>

Version: Accepted Manuscript

Link(s) to article on publisher's website:

<http://dx.doi.org/doi:10.1007/s10801-018-0862-y>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk

The spectra of lifted digraphs ^{*}

C. Dalfó[†], M. A. Fiol[‡], J. Širáň[§]

November 10, 2018

Abstract

We present a method to derive the complete spectrum of the lift Γ^α of a base digraph Γ , with voltage assignments on a (finite) group G . The method is based on assigning to Γ a quotient-like matrix whose entries are elements of the group algebra $\mathbb{C}[G]$, which fully represents Γ^α . This allows us to derive the eigenvectors and eigenvalues of the lift in terms of those of the base digraph and the irreducible characters of G . Thus, our main theorem generalize some previous results of Lovász and Babai concerning the spectra of Cayley digraphs.

Keywords: Digraph, adjacency matrix, regular partition, quotient digraph, spectrum, lifted digraph.

Mathematics Subject Classifications: 05C20; 05C50; 15A18.

^{*}Research of the first two authors is supported by MINECO under project MTM2014-60127-P, and by AGAUR under project 2014SGR1147. The third author acknowledges support from the research grants APVV 0136/12, APVV-15-0220, VEGA 1/0026/16, and VEGA 1/0142/17.

[†]Departament de Matemàtiques, Universitat Politècnica de Catalunya, Barcelona, Catalonia, cristina.dalfo@upc.edu

[‡]Departament de Matemàtiques, Universitat Politècnica de Catalunya; and Barcelona Graduate School of Mathematics, Barcelona, Catalonia, miguel.angel.fiol@upc.edu

[§]Department of Mathematics and Statistics, The Open University, Milton Keynes, UK; and Department of Mathematics and Descriptive Geometry, Slovak University of Technology, Bratislava, Slovak Republic, j.siran@open.ac.uk



The first author has also received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 734922.

1 Preliminaries

As it is well-known, the spectrum of a graph Γ (that is, the eigenvalues of its adjacency matrix) is an important invariant that gives us interesting results about its combinatorial properties, see e.g. the classic textbook of Cvetković, Doob, and Sachs [4]. Thus many efforts have been devoted to derive the spectrum (totally or partially) of some interesting families of graphs. In this framework, Lovász [9] provided a formula which expresses the eigenvalues of Γ in terms of the characters of a transitive subgroup of its automorphism group $\text{Aut } \Gamma$. In the particular case of Cayley graphs (when the automorphism group contains a subgroup G acting regularly on the vertices), Babai [1] derived a more handy formula by different methods. In fact, this formula also holds true for digraphs and arc-colored Cayley graphs. Following these works, here we deal with a more general family of (di)graphs, which are obtained as a type of compounding between a ‘base digraph’ and a Cayley digraph. Our study not only gives the complete spectrum of the obtained digraphs (called ‘lifts’), but also shows how to compute the corresponding eigenvectors.

Through this paper, $\Gamma = (V, E)$ denotes a digraph, with vertex set V and arc set E . An arc from vertex u to vertex v is denoted by either (u, v) , uv , or $u \rightarrow v$. We allow *loops* (that is, arcs from a vertex to itself), and *multiple arcs*. The spectrum of Γ , denoted by $\text{sp } \Gamma = \{\lambda_0^{m_0}, \lambda_1^{m_1}, \dots, \lambda_d^{m_d}\}$, is constituted by the distinct eigenvalues λ_i with the corresponding algebraic multiplicities m_i , for $i \in [0, d]$, of its adjacency matrix \mathbf{A} . (Throughout the paper, for some integer n , we use the notation $[1, n]$ for the set $\{1, \dots, n\}$.)

The paper is organized as follows. In the rest of this section we recall the definition of a voltage digraph and its lift. Then, we introduce a representation of the lifted digraph with a quotient-like matrix whose size equals the order of the (much smaller) base digraph. In particular, it is shown that such a matrix can be used to deduce combinatorial properties of the lifted digraph. Following this approach, and as a main result, Section 2 presents a new method to completely determine the spectrum of the lift by using only the spectrum of the quotient-like matrix and the (irreducible) characters of the group. The results are illustrated by following some examples.

For the concepts and/or results about graphs and digraphs not presented here, we refer the reader to some of the basic textbooks on the subject; for instance, Bang-Jensen and Gutin [2], or Diestel [6].

1.1 Voltage and lifted digraphs

Voltage (di)graphs are, in fact, a type of compounding that consists of connecting together several copies of a (di)graph by setting some (directed) edges between any two copies. More precisely, let Γ be a digraph with vertex set $V = V(\Gamma)$ and arc set $E = E(\Gamma)$. Then, given a group G with generating set Δ , a *voltage assignment* of Γ is a mapping $\alpha : E \rightarrow \Delta$. The

pair (Γ, α) is often called a *voltage digraph*. The *lifted digraph* (or, simply, *lift*) Γ^α is the digraph with vertex set $V(\Gamma^\alpha) = V \times G$ and arc set $E(\Gamma^\alpha) = E \times G$, where there is an arc from vertex (u, g) to vertex (v, h) if and only if $uv \in E$ and $h = g\alpha(uv)$. For example, Figure 1(b) shows the lifted digraph Γ^α for the base digraph $\Gamma = K_2^*$ with voltages shown in Figure 1(a). More precisely, Γ is a complete graph on two vertices $V(\Gamma) = \{a, b\}$, with additional arcs ab, ba, aa, bb , and voltages $\alpha(aa) = \alpha(bb) = \sigma$ and $\alpha(ab) = \alpha(ba) = \rho$, on the group $G = S_3 \cong D_3 = \langle \rho, \sigma \mid \rho^3 = \sigma^2 = (\rho\sigma)^2 = \iota \rangle$. Notice that because of the group role, the symmetry of the obtained lifts usually yields digraphs with large automorphism groups. In particular, when Γ is a singleton with loops, the lift is the Cayley digraph $\text{Cay}(G, \Delta)$. For more information, see the authors' paper [5], or the comprehensive survey of Miller and Širáň [10].

1.2 A matrix representation of the lift

Let us see how we can fully represent a lifted digraph with a matrix whose size equals the order of the base digraph. This approach was initiated by the authors, together with Miller and Ryan, in [5]. Let $\Gamma = (V, E)$ be a digraph with voltage assignment α on the group G . Its *associated matrix* \mathbf{B} is a square matrix indexed by the vertices of Γ , and whose entries are elements of the group algebra $\mathbb{C}[G]$. Namely,

$$(\mathbf{B})_{uv} = \sum_{g \in G} \alpha_g g$$

where

$$\alpha_i = \begin{cases} 1 & \text{if } uv \in E \text{ and } \alpha(uv) = g, \\ 0 & \text{otherwise,} \end{cases}$$

for $i \in [1, n]$. The following result was given in [5].

Lemma 1.1. *Let $(\mathbf{B}^\ell)_{uv} = \sum_{g \in G} a_g^{(\ell)} g$. Then, for every $g, h \in G$, the coefficient $a_g^{(\ell)}$ equals the number of walks of length ℓ in the lifted digraph Γ^α , from vertex (u, h) to vertex (v, hg) . In particular, if $u = v$ and ι denotes the identity element of G , $a_\iota^{(\ell)}$ is the number of walks of length ℓ rooted at every vertex (u, g) , for $g \in G$, of the lift.*

1.3 Some theoretical background

In our study we use representation theory of finite groups. For basics on representation theory and character tables of a group, see for instance, Burrow [3].

We also need to recall the following known result (see e.g. Gould [8]).

Lemma 1.2. *If the power sums $s_k = \sum_{i=1}^d z_i^k$ of some complex numbers z_1, \dots, z_d are known for every $k = 1, \dots, d$, then such numbers are the roots of the monic polynomial*

$$p(z) = \frac{1}{d!} \det \mathbf{C}(z),$$

where $C(z)$ is the following matrix of dimension $d + 1$:

$$C(z) = \begin{pmatrix} z^d & z^{d-1} & z^{d-2} & z^{d-3} & \cdots & z^2 & z & 1 \\ s_1 & 1 & 0 & 0 & \cdots & 0 & 0 & 0 \\ s_2 & s_1 & 2 & 0 & \cdots & 0 & 0 & 0 \\ s_3 & s_2 & s_1 & 3 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ s_{d-1} & s_{d-2} & s_{d-3} & s_{d-4} & \cdots & S_1 & d-1 & 0 \\ s_d & s_{d-1} & s_{d-2} & s_{d-3} & \cdots & s_2 & s_1 & d \end{pmatrix}.$$

2 The spectrum

The following result allows us to compute the spectrum of a lifted digraphs from its associated matrix and the irreducible representations of the group.

Theorem 2.1. *Let $\Gamma = (V, E)$ be a base digraph on r vertices, with a voltage assignment α in a group G with $|G| = n$. Assume that G has ν conjugacy classes with dimensions d_1, \dots, d_ν (so, $\sum_{i=1}^\nu d_i^2 = n$). Let ρ_1, \dots, ρ_ν be the irreducible representations of the group G . Let $\rho_i(\mathbf{B})$ the complex matrix obtained from \mathbf{B} by replacing each $g \in G$ by the $d_i \times d_i$ matrix $\rho_i(g)$, and let $\mu_{u,j}$, $u \in V$, $j \in [1, d_i]$ denote its eigenvalues. Then, the rn eigenvalues of the lift Γ^α are the rd_i eigenvalues of $\rho_i(\mathbf{B})$, for every $i \in [1, \nu]$, each repeated d_i times.*

Proof. Let \mathbf{A} be the adjacency matrix of the lift. First, we prove that, for every $u \in V$ and $j \in [1, d_i]$, every eigenvalue $\mu_{u,j}$ of $\rho_i(\mathbf{B})$ gives rise to d_i (equal) eigenvalue of \mathbf{A} . To this end, let \mathbf{U}_i the $rd_i \times rd_i$ matrix whose columns are the eigenvectors of $\rho_i(\mathbf{B})$. Let \mathbf{D}_i be the diagonal matrix having such eigenvalues as its entries. For every $u \in V$, let \mathbf{x}_u be the $d_i \times rd_i$ submatrix of \mathbf{U}_i having files indexed with (u, j) , $j \in [1, d_i]$. Then, from $\rho_i(\mathbf{B})\mathbf{U}_i = \mathbf{U}_i\mathbf{D}_i$ we have

$$\sum_{uv \in E} \rho_i(\alpha(uv))\mathbf{x}_v = \mathbf{x}_u\mathbf{D}_i \quad \text{for } u \in V. \quad (1)$$

Now for each vertex (u, g) of Γ^α , consider the $d_i \times rd_i$ matrix

$$\phi_{(u,g)} = \rho_i(g)\mathbf{x}_u.$$

Then,

$$\begin{aligned} \phi_{(u,g)}\mathbf{D}_i &= \rho_i(g)\mathbf{x}_u\mathbf{D}_i = \rho_i(g) \sum_{uv \in E} \rho_i(\alpha(uv))\mathbf{x}_v \\ &= \sum_{uv \in E} \rho_i(g\alpha(uv))\mathbf{x}_v = \sum_{uv \in E} \phi_{(v,g\alpha(uv))}. \end{aligned}$$

But this means that, for every pair of fixed elements $k \in [1, d_i]$ and $(u, j) \in \{V \times [1, d_i]\}$, the vector obtained by taking the $(k, (u, j))$ -entry of every matrix $\phi_{(v, h)}$, for $(v, h) \in V(\Gamma^\alpha)$, is an eigenvector of the lift Γ^α with eigenvalue $\mu_{u, j}$. Since there are d_i possible choices for k , the same holds for the eigenvectors of $\mu_{u, j}$.

Moreover, by Lemma 1.1, if $(\mathbf{B}^\ell)_{uu} = \sum_{g \in G} a_{(u, g)}^{(\ell)} g$, the total number of rooted closed ℓ -walks in Γ^α is

$$\text{tr}(\mathbf{A}^\ell) = \sum_{\lambda \in \text{sp } \Gamma^\alpha} \lambda^\ell = n \sum_{u \in V} a_{(u, \iota)}^{(\ell)}$$

since, in the lift, the number of (u, g) -rooted closed ℓ -walks does not depend on $g \in G$. Moreover, by the ‘Great Orthogonality Theorem’ (see, e.g. [3]), we have $\sum_{g \in G} \rho_i(g) = \mathbf{0}$ for every $i \neq 1$ (with ρ_1 being the trivial representation). Then,

$$a_{(u, \iota)}^{(\ell)} = \frac{1}{n} \sum_{i=1}^{\nu} d_i (\rho_i(\mathbf{B}^\ell))_{uu}$$

and, hence,

$$\sum_{\lambda \in \text{sp } \Gamma^\alpha} \lambda^\ell = \text{tr}(\mathbf{A}^\ell) = \sum_{i=1}^{\nu} d_i \text{tr}(\rho_i(\mathbf{B}^\ell)) = \sum_{i=1}^{\nu} \sum_{\mu \in \text{sp } \rho_i(\mathbf{B}^\ell)} d_i \mu^\ell.$$

(Note that, in the sum on the right, we have $r \sum_{i=1}^{\nu} d_i^2 = rn$ terms, which is the number of eigenvalues of the adjacency matrix \mathbf{A} of the lift Γ^α .) As the above equality holds for every $\ell = 1, 2, \dots$, by Lemma 1.2 both (multi)sets of eigenvalues must coincide. \square

As a consequence, we have the following result in terms of the (irreducible) characters $\chi_i(g)$, for $g \in G$, of the group. For stating it, let us introduce some additional notation: Let P_ℓ be the set of closed walks of length $\ell \geq 1$ in Γ . If $p \in P_\ell$, say $u_0 \rightarrow u_1 \rightarrow \dots \rightarrow u_{\ell-1} \rightarrow u_\ell (= u_0)$, let

$$\chi_i(p) = \prod_{j=0}^{\ell-1} \chi_i(\alpha(u_j u_{j+1})).$$

Corollary 2.2. *Using the same notation as above, for each $i \in [1, \nu]$, the eigenvalues $\lambda_{u, j}$, for $u \in V$ and $j \in [1, d_i]$, of the lift Γ^α , are the solutions (each repeated d_i times) of the system*

$$\sum_{u \in V, j \in [1, d_i]} \lambda_{u, j}^\ell = \sum_{p \in P_\ell} \chi_i(p), \quad \ell = 1, \dots, rd_i. \quad (2)$$

Proof. By Theorem 2.1, the above left sum of the powers is $\text{tr}(\rho_i(\mathbf{B})^\ell)$, whereas the right expression corresponds to $\chi_i(\text{tr}(\mathbf{B}^\ell))$ (where the ℓ -power of \mathbf{B} and its trace is computed in $\mathbb{C}[G]$, and $\chi_i(\sum_{g \in G} a_g g) = \sum_{g \in G} a_g \chi_i(g)$). Then, the result follows from

$$\text{tr}(\rho_i(\mathbf{B})^\ell) = \sum_{u \in V} \text{tr}[\rho_i((\mathbf{B}^\ell)_{uu})] = \sum_{u \in V} \chi_i((\mathbf{B}^\ell)_{uu}) = \text{tr}(\chi_i(\mathbf{B}^\ell)).$$

□

Notice that, by Lemma 1.2, the equalities in (2) lead to a polynomial of degree rd_i , with roots the required eigenvalues $\lambda_{u,j}$.

As commented in Section 1.1, when Γ consists of one vertex with loops, then Γ^α is a Cayley digraph and (2) gives the result of Babai [1]. Another extreme case is when $d_i = 1$ for some i . Then, we simply have $\lambda_{u,1} = \mu_{u,1}$ for every $u \in V$. For instance, when G is Abelian, $\nu = n$, and this holds for every $i \in [1, n]$. This case was dealt with by the authors, Miller, and Ryan in [5].

2.1 An example

Let us consider again the lift described in Subsection 1.1 and shown Figure 1. Then, the base graph K_2^* has voltages on the symmetric group $S_3 \cong D_3 = \langle \rho, \sigma : \rho^3 = \sigma^2 = (\rho\sigma)^2 = \iota \rangle$, with characters shown in Table 1, and associated matrix

$$\mathbf{B} = \begin{pmatrix} \sigma & \iota + \rho \\ \iota + \rho & \sigma \end{pmatrix}.$$

Note that the edge (two opposite arcs forming a digon) of the base digraph gives rise to the entries ι 's for the voltages assigned to the corresponding arcs.

$S_3 \setminus g$	ι	$\sigma, \sigma\rho, \sigma\rho^2$	ρ, ρ^2
$\chi_1 (d_1 = 1)$	1	1	1
$\chi_2 (d_2 = 1)$	1	-1	1
$\chi_3 (d_3 = 2)$	2	0	-1

Table 1: The character table of the symmetric group S_3 .

The obtained lifted digraph Γ^α has spectrum

$$\text{sp } \Gamma^\alpha = \{3^{(1)}, 1^{(3)}, 0^{(4)}, -1^{(3)}, -3^{(1)}\}. \quad (3)$$

Then, we get the complex matrices

$$\chi_1(\mathbf{B}) = \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}, \quad \chi_2(\mathbf{B}) = \begin{pmatrix} -1 & 2 \\ 2 & -1 \end{pmatrix}, \quad \chi_3(\mathbf{B}) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

Then, by Corollary 2.2:

- χ_1 : Since $d_1 = 1$, two eigenvalues of Γ^α are

$$\{3, -1\} = \text{ev } \chi_1(\mathbf{B}).$$




Figure 1: The base digraph K_2^* , on the group S_3 , and its lift

- χ_2 : Since $d_2 = 1$, two eigenvalues of Γ^α are

$$\{-3, 1\} = \text{ev } \chi_2(\mathbf{B}).$$

- χ_3 : Since $d_3 = 2$, we consider all the possible closed walks of lengths $\ell = 1, 2, 3, 4$ in \mathbf{B} , which gives the system

$$\begin{aligned} \lambda_{u,0} + \lambda_{u,1} + \lambda_{v,0} + \lambda_{v,1} &= 0 \\ \lambda_{u,0}^2 + \lambda_{u,1}^2 + \lambda_{v,0}^2 + \lambda_{v,1}^2 &= 2 \\ \lambda_{u,0}^3 + \lambda_{u,1}^3 + \lambda_{v,0}^3 + \lambda_{v,1}^3 &= 0 \\ \lambda_{u,0}^4 + \lambda_{u,1}^4 + \lambda_{v,0}^4 + \lambda_{v,1}^4 &= 2, \end{aligned}$$

with solutions $1, 0, 0, -1$

Then, as these last eigenvalues have to be considered twice, this completes the eigenvalue multiset of Γ^α , in agreement with (3).

References

- [1] L. Babai, Spectra of Cayley graphs, *J. Combin. Theory Ser. B* **27** (1979) 180–189.
- [2] J. Bang-Jensen and G. Gutin, *Digraphs. Theory, Algorithms and Applications*. Second edition. Springer Monographs in Mathematics. Springer-Verlag London, Ltd., London, 2009.
- [3] M. Burrow, *Representation Theory of Finite Groups*, Dover Publications, Inc., New York, 1993.

- [4] D. Cvetković, M. Doob, and H. Sachs, *Spectra of Graphs. Theory and Applications* 3rd edition, Johann Ambrosius Barth, Heidelberg, 1995.
- [5] C. Dalfó, M. A. Fiol, M. Miller, J. Ryan, and J. Širáň, An algebraic approach to lifts of digraphs, submitted (2017), [arXiv:1612.08855v2 \[math.CO\]](#).
- [6] R. Diestel, *Graph Theory* (4th ed.), Graduate Texts in Mathematics **173**, Springer-Verlag, Heilderberg, 2010.
- [7] R. Gera and P. Stănică, The spectrum of generalized Petersen graphs, *Australasian J. Combin.* **49** (2011) 39–45.
- [8] H. W. Gould, The Girard-Waring power sum formulas for symmetric functions and Fibonacci sequences, *Fibonacci Quart.* **37** (1999), no. 2, 135–140.
- [9] L. Lovász, Spectra of graphs with transitive groups, *Period. Math. Hungar.* **6** (1975) 191–196.
- [10] M. Miller and J. Širáň, Moore graphs and beyond: A survey of the degree/diameter problem, *Electron. J. Combin.* **20(2)** (2013) #DS14v2.